

Relations between the leading terms of a polynomial automorphism

Philippe Bonnet and Stéphane Vénéreau
Mathematisches Institut, Universität Basel
Rheinsprung 21, CH-4051 Basel, Switzerland
E-mails: philippe.bonnet@unibas.ch,
stephane.venereau@unibas.ch

Abstract

Let I be the ideal of relations between the leading terms of the polynomials defining an automorphism of K^n . In this paper, we prove the existence of a locally nilpotent derivation which preserves I . Moreover, if I is principal, i.e. $I = (R)$, we compute an upper bound for $\deg_2(R)$ for some degree function $\deg_2$ defined by the automorphism. As applications, we determine all the principal ideals of relations for automorphisms of K^3 and deduce two elementary proofs of the Jung-van der Kulk Theorem about the tameness of automorphisms of K^2 .

1 Introduction

Throughout this paper, K denotes an algebraically closed field of characteristic zero. For any positive real numbers $w_1, \dots, w_n$, $\deg_1$ stands for the weighted homogeneous degree on the ring $K[x_1, \dots, x_n]$ of polynomials in n variables, which assigns the weights w_i to every x_i . We say that $\deg_1$ defines a positive weighted homogeneous degree (in short "p.w.h. "). When all the weights are equal to 1, we speak of the (standard) homogeneous degree on $K[x_1, \dots, x_n]$. Given any polynomial $f = \sum a_{k_1, \dots, k_n} x_1^{k_1} \cdots x_n^{k_n}$ of degree d for $\deg_1$, its leading term $\bar{f}$ (with respect to $\deg_1$) is defined as:

$$\bar{f} = \sum_{k_1 w_1 + \dots + k_n w_n = d} a_{k_1, \dots, k_n} x_1^{k_1} \cdots x_n^{k_n}$$

In order to understand polynomial automorphisms, it is natural to study the leading terms of their components. More precisely, let $\Phi = (f_1, \dots, f_n)$ be a polynomial automorphism of K^n . What can be said about $\bar{f}_1, \dots, \bar{f}_n$? Except for $n \leq 2$ (see section 5 below), nothing is known. In this paper, we would like to determine the algebraic relations between these leading terms.

Definition 1.1 *Given a p.w.h. degree on $K[x_1, \dots, x_n]$ and an automorphism $(f_1, \dots, f_n)$ of K^n , the set I of polynomials $P \in K[x_1, \dots, x_n]$ such that $P(\bar{f}_1, \dots, \bar{f}_n) = 0$ is a prime ideal called the ideal of relations.*

Given a p.w.h. degree $\deg_1$ on $K[x_1, \dots, x_n]$ and an automorphism $(f_1, \dots, f_n)$ of K^n , we introduce a new p.w.h. degree $\deg_2$ on $K[x_1, \dots, x_n]$, which assigns the weight $d_i = \deg_1(f_i)$

to every variable x_i . Recall that a K -derivation ∂ of a K -algebra A is locally nilpotent if, for any $f \in A$, there exists an order $k \geq 0$ with $\partial^k(f) = 0$. Such a derivation is called an *LND*. After some simple observations in section 2, we will establish, in sections 3 and 4, the following two properties for ideals of relations:

Theorem 1.2 *Let $\Phi = (f_1, \dots, f_n)$ be an automorphism of K^n , let $\deg_1$ be a p.w.h. degree on $K[x_1, \dots, x_n]$ and let I be the ideal of relations. Then, there exists a nonzero locally nilpotent derivation ∂ of $K[x_1, \dots, x_n]$ such that $\partial(I) \subset I$. Moreover, if this ideal is principal, i.e. $I = (R)$, then $\partial(R) = 0$.*

Theorem 1.3 *Let $\Phi = (f_1, \dots, f_n)$ be an automorphism of K^n . Assume that $\deg_1$ is the standard homogeneous degree on $K[x_1, \dots, x_n]$. If the ideal of relations is principal, i.e. $I = (R)$, then $\deg_2(R) \leq d_1 + \dots + d_n - n + 1$.*

Recall that an automorphism $\tau : K^n \rightarrow K^n$ is elementary if, up to a permutation of $x_1, \dots, x_n$, it can be written as $\tau(x_1, \dots, x_n) = (x_1, \dots, x_n + P)$, where $P \in K[x_1, \dots, x_{n-1}]$. An automorphism of K^n is *tame* if it is a composition of affine and elementary automorphisms. The Jung-van der Kulk Theorem states that *every automorphism of K^2 is tame* (see [Jun, vdK]). In section 5, we give two elementary proofs of this result, based on Theorems 1.2 and 1.3.

However the situation for $n \geq 3$ becomes more complicated. Indeed, there exist nontame automorphisms in dimension 3 (see [SU2]). In sections 6, 7 and 8, applying Theorems 1.2 and 1.3, we can determine all possible principal ideals of relations when $n = 3$, thereby proving:

Theorem 1.4 *Let $\Phi = (f_1, f_2, f_3)$ be an automorphism of K^3 , and let $\deg_1$ be the standard homogeneous degree on $K[x_1, x_2, x_3]$. Up to a permutation of the f_i , we may assume that $d_1 \leq d_2 \leq d_3$. Assume that the ideal I of relations is principal, i.e. $I = (R)$. Then there exists a $\deg_2$ -homogeneous polynomial $h(x_1, x_2)$ whose $\deg_2$ is equal to $\deg_2(x_3) = \deg_1(f_3)$ or $-\infty$ (h might be zero) such that R is proportional to one of the following polynomials, where $x'_3 := x_3 + h(x_1, x_2)$:*

- | | | |
|------|---|--|
| (1) | 0 | (affine case); |
| (2) | x'_3 | ("elementarily reducible" case); |
| (3) | $x_1^{e_1} + cx_2^{e_2}$ | with $c \in K^*$, $\gcd(e_1, e_2) = 1$; |
| (4) | $(x_2 + ax_1^{e_1})x'_3 + cx_1^k$ | with $c \in K^*$, $a \in K$, $k \geq 2$, $e_1 \geq 1$; |
| (5) | $x_1^k x'_3 + P(x_1, x_2)$ | with $k \geq 1$, P $\deg_2$ -homogeneous ; |
| (6) | $x_3'^2 + cx_2^3$ | with $c \in K^*$; |
| (7) | $x_3'^2 + cx_1 x_2^2$ | with $c \in K^*$; |
| (8) | $x_3'^2 + cx_1^{r_1}$ | with $c \in K^*$, $r_1 \in 3 + 2\mathbb{N}$; |
| (9) | $x_3'^2 + cx_1^{r_1} x_2$ | with $c \in K^*$, $r_1 \geq 1$; |
| (10) | $x_3'^2 + ax_1^{e_1} + bx_2^2$ | with $ab \in K^*$, $e_1 \in 3 + 2\mathbb{N}$; |
| (11) | $x_3'^2 + (ax_1^{e_1} + bx_2)x_1^{r_1}$ | with $ab \in K^*$, $e_1 \geq 1$; |
| (12) | $x_3'^2 + (a_1 x_1^{e_1} + b_1 x_2)(a_2 x_1^{e_1} + b_2 x_2)$ | with $a_1 b_2 - b_1 a_2 \neq 0$, $e_1 \geq 1$; |
| (13) | $x_3'^2 + (a_1 x_1 + b_1 x_2)(a_2 x_1 + b_2 x_2)^2$ | with $(a_i, b_i) \neq (0, 0)$ for $i = 1, 2$; |
| (14) | $x_3'^2 + (ax_1^{e_1} + bx_2)^2 x_1$ | with $ab \neq 0$, $e_1 \geq 2$. |

Two polynomials f and g of $K[x_1, \dots, x_n]$ are said *tamely equivalent* if there exists a tame automorphism ψ of K^n such that $f = g \circ \psi$. With x'_3 as in the theorem, $R(x_1, x_2, x_3)$ and $R(x_1, x_2, x'_3)$ are of course always tamely equivalent.

As a consequence of Theorem 1.4, we will prove in section 9 the following

Corollary 1.5 *Let Φ be an automorphism of K^3 and let $\deg_1$ be the standard homogeneous degree on $K[x_1, x_2, x_3]$. If the ideal I of relations is principal, i.e. $I = (R)$, then R is tamely equivalent to one of the following polynomials: 0 , x_3 , $x_1^r + x_2^s$ with $\gcd(r, s) = 1$, $x_1^k x_3 + P(x_1, x_2)$ with $k \geq 1$.*

2 Some simple but useful facts

A derivation ∂ on a general commutative ring A with the same local nilpotence condition as in the introduction is also called a LND. One may define a function $\deg_\partial$ by $\deg_\partial(a) = \max\{n \mid \partial^n(a) \neq 0\}$ if $a \neq 0$ and $\deg_\partial(0) = -\infty$. The following lemma is well-known, see e.g. [ML1]:

Lemma 2.1 *If A is a domain then $\deg_\partial$ is a degree function, i.e. $\deg_\partial(a+b) \leq \max\{\deg_\partial(a), \deg_\partial(b)\}$ and $\deg_\partial(ab) = \deg_\partial(a) + \deg_\partial(b)$.*

Corollary 2.2 *If A is a domain, $a \in A$ and $\partial(a) = ba$ for some $b \in A$ then $\partial(a) = ba = 0$.*

Proof: If $\partial(a) = ba \neq 0$ then $\deg_\partial(a) = \deg_\partial(\partial a) + 1 = \deg_\partial(ab) + 1 = \deg_\partial(a) + \deg_\partial(b) + 1$ and one gets $\deg_\partial(b) = -1$ which is impossible. ■

We will also need to consider the *jacobian* (determinant) of n polynomials $P_1, \dots, P_n \in K[x_1, \dots, x_n]$:

$$j(P_1, \dots, P_n) = \det(\partial P_i / \partial x_j)_{1 \leq i, j \leq n}.$$

It is clearly a K -derivation in every one of its entries i.e. every $P_i \mapsto j(P_1, \dots, P_n)$ is a derivation (when all the other P_j 's are fixed). From its definition, it is also clear that $\deg_1(j(P_1, \dots, P_n)) \leq \deg_1(P_1) + \dots + \deg_1(P_n) - (\deg_1(x_1) + \dots + \deg_1(x_n))$. In particular, if $\deg_1$ is the standard homogeneous degree then

$$(15) \quad \deg_1(j(P_1, \dots, P_n)) \leq \deg_1(P_1) + \dots + \deg_1(P_n) - n.$$

Note also that, if $\Psi = (h_1, \dots, h_n)$ is an automorphism, then $j(h_1, \dots, h_n) = \mu \in K^*$ (well-known fact, consequence of the chain rule applied on its composition with its inverse) and one has, $\forall i = 1, \dots, n$,

$$\begin{aligned} j(h_1, \dots, h_{i-1}, P \circ \Psi, h_{i+1}, \dots, h_n) &= j(h_1, \dots, h_{i-1}, P(h_1, \dots, h_n), h_{i+1}, \dots, h_n) \\ &= j(h_1, \dots, h_n) \frac{\partial P}{\partial x_i}(h_1, \dots, h_n) \\ (16) \quad j(h_1, \dots, h_{i-1}, P \circ \Psi, h_{i+1}, \dots, h_n) &= \mu \frac{\partial P}{\partial x_i} \circ \Psi. \end{aligned}$$

Now let $\deg_1$, $\Phi = (f_1, \dots, f_n)$, I and $\deg_2$ be as in the introduction. Recall that $\bar{f}$ denotes the leading term of the polynomial $f \in K[x_1, \dots, x_n]$ with respect to $\deg_1$. Likewise, we will denote $\tilde{P}$ the leading term of a polynomial $P \in K[x_1, \dots, x_n]$ with respect to $\deg_2$.

Lemma 2.3 *Let P be any polynomial in $K[x_1, \dots, x_n]$. Then $\deg_1(P \circ \Phi) \leq \deg_2(P)$ and the inequality is strict if and only if $\tilde{P} \in I$ and $P \neq 0$.*

Proof: If $P = 0$ this is trivial so let's us assume this is not the case.

If $p_{i_1 \dots i_n}$ denotes the coefficient of $x_1^{i_1} \dots x_n^{i_n}$ in P then, by definition of the degree $\deg_2$, there exists an index $i_1 \dots i_n$ such that $i_1 d_1 + \dots + i_n d_n = \deg_2(P)$ and $p_{i_1 \dots i_n} \neq 0$ and one may write:

$$P(x_1, \dots, x_n) = \sum_{i_1 d_1 + \dots + i_n d_n = \deg_2(P)} p_{i_1 \dots i_n} x_1^{i_1} \dots x_n^{i_n} + \sum_{i_1 d_1 + \dots + i_n d_n < \deg_2(P)} p_{i_1 \dots i_n} x_1^{i_1} \dots x_n^{i_n}$$

and, by definition of $\tilde{P}$,

$$P(x_1, \dots, x_n) = \tilde{P}(x_1, \dots, x_n) + \sum_{i_1 d_1 + \dots + i_n d_n < \deg_2(P)} p_{i_1 \dots i_n} x_1^{i_1} \dots x_n^{i_n}$$

Composing with $\Phi = (f_1, \dots, f_n)$ we get

$$\begin{aligned} P \circ \phi = P(f_1, \dots, f_n) &= \tilde{P}(f_1, \dots, f_n) + \sum_{i_1 d_1 + \dots + i_n d_n < \deg_2(P)} p_{i_1 \dots i_n} f_1^{i_1} \dots f_n^{i_n} \\ &= \tilde{P}(\bar{f}_1, \dots, \bar{f}_n) + ldt \end{aligned}$$

with

$$ldt = \tilde{P}(f_1, \dots, f_n) - \tilde{P}(\bar{f}_1, \dots, \bar{f}_n) + \sum_{i_1 d_1 + \dots + i_n d_n < \deg_2(P)} p_{i_1 \dots i_n} f_1^{i_1} \dots f_n^{i_n}.$$

One has

$$\deg_1(ldt) \leq \max\{\deg_1(\tilde{P}(f_1, \dots, f_n) - \tilde{P}(\bar{f}_1, \dots, \bar{f}_n)), \deg_1\left(\sum_{i_1 d_1 + \dots + i_n d_n < \deg_2(P)} p_{i_1 \dots i_n} f_1^{i_1} \dots f_n^{i_n}\right)\}$$

hence $\deg_1(ldt) < \deg_2(P)$. Moreover either $\deg_1(\tilde{P}(\bar{f}_1, \dots, \bar{f}_n)) = \deg_2(P)$ or $\tilde{P}(\bar{f}_1, \dots, \bar{f}_n) = 0$.

Now let's sum up:

$$\begin{cases} P \circ \Phi = \tilde{P}(\bar{f}_1, \dots, \bar{f}_n) + ldt \\ \deg_1(\tilde{P}(\bar{f}_1, \dots, \bar{f}_n)) = \deg_2(P) > \deg_1(ldt) \text{ or } \tilde{P}(\bar{f}_1, \dots, \bar{f}_n) = 0 \end{cases}$$

and the result follows from the following trivial remark (which holds for any degree function):

$$\deg a + b \leq \max\{\deg a, \deg b\} \quad \text{and} \quad \deg a < \deg b \Rightarrow \deg a + b = \deg b$$

applied to $\deg = \deg_1$, $a = \tilde{P}(\bar{f}_1, \dots, \bar{f}_n)$ and $b = ldt$.

■

Note that, in the case when $\deg_1$ is the standard homogeneous degree, the weights of $\deg_2$, namely the $d_i = \deg_1(f_i)$'s are all at least 1, since $\Phi = (f_1, \dots, f_n)$ is an automorphism. Consequently one gets the implication: $\deg_1(P) > 1 \Rightarrow \deg_2(P) > 1$.

Corollary 2.4 *If $\deg_1$ is the standard homogeneous degree and $P \in K[x_1, \dots, x_n]$ is such that $\deg_1(P \circ \Phi) = 1$ then $\tilde{P} \in I$ unless P is affine i.e. $\deg_1(P) \leq 1$.*

Proof: If P is not affine then $\deg_2(P) > 1 = \deg_1(P \circ \Phi)$ and it follows from Lemma 2.3 that $\tilde{P} \in I$. ■

Corollary 2.5 *If $\deg_1$ is the standard homogeneous degree then $I = (0)$ if and only if Φ is an affine automorphism i.e. $\deg_1(f_i) = 1, \forall i = 1, \dots, n$.*

Proof: If Φ is affine then

$$\bar{\Phi} := (\bar{f}_1, \dots, \bar{f}_n) = (f_1 - f_1(0, \dots, 0), \dots, f_n - f_n(0, \dots, 0)) = \tau \circ \Phi$$

where τ is the automorphism (translation) $(x_1 - f_1(0, \dots, 0), \dots, x_n - f_n(0, \dots, 0))$ also $\bar{\Phi}$ itself is an automorphism preventing that $P \circ \bar{\Phi} = P(\bar{f}_1, \dots, \bar{f}_n) = 0$ for any $P \neq 0$. It follows that $I = (0)$.

Assume now that $I = (0)$. Let $\Phi^{-1} = (g_1, \dots, g_n)$ be the inverse of Φ . One has $\forall i = 1, \dots, n$, $g_i \circ \Phi = x_i$ hence $1 = \deg_1(x_i) = \deg_1(g_i \circ \Phi)$ and $0 \neq g_i \notin I$. By Corollary 2.4, g_i is affine $\forall i = 1, \dots, n$, hence so are $\Phi^{-1} = (g_1, \dots, g_n)$ and Φ . ■

Remark that for p.w.h. degrees other than the standard one, none of the implications in Corollary 2.5 holds. Indeed, take the weights $w_1 = \deg_1(x_1) = 1, w_2 = \deg_1(x_2) = 3$. Then $(f_1, f_2) = (x_1 + x_2, x_2)$ is an affine automorphism but $\bar{f}_1 = x_2$ and $\bar{f}_2 = x_2$ fullfill the non trivial relation $\bar{f}_1 - \bar{f}_2 = 0$.

Conversely $(f_1, f_2) = (x_1, x_2 + x_1^2)$ is a non-affine automorphism but $\bar{f}_1 = x_1$ and $\bar{f}_2 = x_2$ are algebraically independant.

Finally, we point out an easy fact linking degree and partial derivative (here $\frac{\partial}{\partial x_n}$) in the following

Lemma 2.6 *Let P be a polynomial in $K[x_1, \dots, x_n]$. If $\frac{\partial \tilde{P}}{\partial x_n} \neq 0$ then $\widetilde{\frac{\partial P}{\partial x_n}} = \frac{\partial \tilde{P}}{\partial x_n}$ and $\deg_2(\frac{\partial P}{\partial x_n}) = \deg_2(P) - d_n$. More generally, while $\frac{\partial^k \tilde{P}}{\partial x_n^k} \neq 0$ one has $\widetilde{\frac{\partial^k P}{\partial x_n^k}} = \frac{\partial^k \tilde{P}}{\partial x_n^k}$ and $\deg_2(\frac{\partial^k P}{\partial x_n^k}) = \deg_2(P) - kd_n$.*

Proof: Derivating $P = \tilde{P} + ldt$ where $\deg_2(ldt) < \deg_2(P) = \deg_2(\tilde{P})$ one gets

$$\frac{\partial P}{\partial x_n} = \frac{\partial \tilde{P}}{\partial x_n} + \frac{\partial ldt}{\partial x_n}.$$

Since $\tilde{P}$ is $\deg_2$ -homogeneous, so is $\frac{\partial \tilde{P}}{\partial x_n}$ and its $\deg_2$ is equal to $\deg_2(P) - \deg_2(x_n) = \deg_2(P) - d_n$. On the other hand, one has $\deg_2(\frac{\partial ldt}{\partial x_n}) \leq \deg_2(ldt) - d_n < \deg_2(P) - d_n$ and everything comes from the equality above. ■

3 Proof of the first Theorem

Let ∂ be a K -derivation of $K[x_1, \dots, x_n]$. Given a p.w.h. degree $\deg$ on $K[x_1, \dots, x_n]$, the degree of ∂ is defined as:

$$\deg(\partial) = \sup\{\deg(\partial(P)) - \deg(P) \mid P \in K[x_1, \dots, x_n] - \{0\}\}$$

Note that this degree is upper-bounded, i.e. $\deg(\partial) < +\infty$, and equals $-\infty$ if and only if ∂ is the zero derivation. To see this, it suffices to check that:

$$\deg(\partial) = \sup\{\deg(\partial(x_i)) - \deg(x_i) \mid i = 1, \dots, n\}$$

By construction, we have the following inequality:

$$\forall P \in K[x_1, \dots, x_n], \quad \deg(\partial(P)) \leq \deg(\partial) + \deg(P).$$

The leading term $\bar{\partial}$ of a k -derivation ∂ is the k -linear endomorphism of $K[x_1, \dots, x_n]$, which maps every weighted homogeneous polynomial P to the weighted homogeneous term of $\partial(P)$ of degree $\deg(\partial) + \deg(P)$. By construction, $\bar{\partial}$ is a k -derivation of $K[x_1, \dots, x_n]$. More precisely, assume that $\partial = \sum_i a_i \partial / \partial x_i$, where every a_i is a polynomial. If r is the degree of ∂ and if b_i denotes the weighted homogeneous term of degree $r + \deg(x_i)$ of a_i , then:

$$\bar{\partial} = \sum_{i=1}^n b_i \frac{\partial}{\partial x_i}.$$

Let $\Phi = (f_1, \dots, f_n)$ be a polynomial automorphism of K^n , with jacobian equal to $\lambda \in k$, and denote by $\Phi^{-1} = (g_1, \dots, g_n)$ its inverse (also with jacobian λ^{-1}). Let $\deg_1$ be a p.w.h. degree, which assigns the weight w_i to each variable x_i , and let $\deg_2$ be the degree defined in the introduction. For any $i = 1, \dots, n$, consider the k -derivations δ_i and Δ_i on $K[x_1, \dots, x_n]$, defined for any polynomial P by the formulas:

$$\delta_i(P) = \lambda^{-1} \frac{\partial P}{\partial x_i} \quad \text{and} \quad \Delta_i(P) = j(g_1, \dots, g_{i-1}, P, g_{i+1}, \dots, g_n).$$

Note that each derivation δ_i has degree $-w_i$ with respect to $\deg_1$. Theorem 1.2 will be a straightforward consequence of the following lemmas, where we prove that the leading term of one of the Δ_i for $\deg_2$ is an LND which stabilizes the ideal I of relations.

Lemma 3.1 *For any index i of $\{1, \dots, n\}$ and any polynomial P of $K[x_1, \dots, x_n]$, we have $\Delta_i(P \circ \Phi^{-1}) = \delta_i(P) \circ \Phi^{-1}$ and $\Delta_i(P) \circ \Phi = \delta_i(P \circ \Phi)$. Moreover, every derivation Δ_i is locally nilpotent.*

Proof: For any index $i = 1, \dots, n$ and any polynomial P of $K[x_1, \dots, x_n]$ we have, by equality (16):

$$\Delta_i(P \circ \Phi^{-1}) = j(g_1, \dots, g_{i-1}, P \circ \Phi^{-1}, g_{i+1}, \dots, g_n) = \lambda^{-1} \frac{\partial P}{\partial x_i} \circ \Phi^{-1} = \delta_i(P) \circ \Phi^{-1}.$$

Now, for any polynomial P , if we set $Q = P \circ \Phi$, then we find:

$$\Delta_i(P) \circ \Phi = \Delta_i(Q \circ \Phi^{-1}) \circ \Phi = \delta_i(Q) \circ \Phi^{-1} \circ \Phi = \delta_i(Q) = \delta_i(P \circ \Phi).$$

Using the second formula, one can prove by induction on $k \geq 1$ that $\Delta_i^k(P) \circ \Phi = \delta_i^k(P \circ \Phi)$ for any P . Now fix a polynomial P . Since δ_i is locally nilpotent, there exists an order $r \geq 1$ such that $\delta_i^r(P \circ \Phi) = 0$. In particular $\Delta_i^r(P) \circ \Phi = 0$. Since Φ is an automorphism, $\Delta_i^r(P) = 0$. Since this holds for any P , Δ_i is locally nilpotent. ■

Lemma 3.2 *There exists an index i such that $\deg_2(\Delta_i) \geq \deg_1(\delta_i) = -w_i$.*

Proof: Suppose on the contrary that $\deg_2(\Delta_i) < \deg_1(\delta_i) = -w_i$ for any $i = 1, \dots, n$. Fix an index j for which $d_j = \min_i \{d_i\}$. Using Lemma 3.1 and the fact that $x_j = f_j \circ \Phi^{-1}$, we obtain for any index i :

$$\deg_2(\Delta_i(x_j)) = \deg_2(\Delta_i(f_j \circ \Phi^{-1})) = \deg_2(\delta_i(f_j) \circ \Phi^{-1}) < d_j - w_i$$

Since d_j is minimal among all the d_i , every non-constant polynomial has degree $\geq d_j$ with respect to $\deg_2$. Since every weight w_i is positive, $d_j - w_i < d_j$ for any i . So $\delta_i(f_j) \circ \Phi^{-1}$ is constant for any i . Since Φ is an automorphism, $\delta_i(f_j)$ is constant for any i . This implies there exists some constants $a, a_1, \dots, a_n$ such that:

$$f_j = a + a_1 x_1 + \dots + a_n x_n$$

In particular, d_j is the maximum of the w_i for which $a_i \neq 0$. Now fix an index l such that $d_j = w_l$ and $a_l \neq 0$. As before, we obtain:

$$\deg_2(\Delta_l(x_j)) = \deg_2(\delta_l(f_j) \circ \Phi^{-1}) < d_j - w_l = 0$$

Since every weight w_i is positive, the polynomial $\delta_l(f_j) \circ \Phi^{-1}$ must be equal to zero. But this is impossible since $\delta_l(f_j) = a_l \neq 0$. ■

Lemma 3.3 *Let i be an index such that $\deg_2(\Delta_i) \geq -w_i$. Then the leading part $\overline{\Delta_i}$ of Δ_i with respect to $\deg_2$ is locally nilpotent and stabilizes the ideal I .*

Proof: Let i be an index such that $\deg_2(\Delta_i) \geq -w_i = \deg_1(\delta_i)$. Since Δ_i is locally nilpotent by Lemma 6.1, its leading term $\overline{\Delta_i}$ is also locally nilpotent (see [vdE]). So we only need to prove that $\overline{\Delta_i}$ stabilizes I . Since the $\overline{f_i}$ are $\deg_1$ -homogeneous of degrees $d_1, \dots, d_n$, I is a weighted homogeneous ideal for $\deg_2$. Let P be any nonzero $\deg_2$ -homogeneous element of I . Using successively Lemma 2.3, the definition of the degree of a derivation and Lemma 3.1, we find:

$$\begin{aligned} \deg_2(\Delta_i) + \deg_2(P) &> \deg_1(\delta_i) + \deg_1(P \circ \Phi) \\ &> \deg_1(\delta_i(P \circ \Phi)) \\ &> \deg_1(\Delta_i(P) \circ \Phi) \end{aligned}$$

If $\deg_2(\Delta_i(P)) < \deg_2(\Delta_i) + \deg_2(P)$, then $\overline{\Delta_i}(P) = 0$ and there is nothing to prove. So assume that $\deg_2(\Delta_i(P)) = \deg_2(\Delta_i) + \deg_2(P)$. By the inequality proved above, we get:

$$\deg_2(\Delta_i(P)) > \deg_1(\Delta_i(P) \circ \Phi)$$

Therefore, $\overline{\Delta_i}(P)$ belongs to I by Lemma 2.3. Since $\deg_2(\Delta_i(P)) = \deg_2(\Delta_i) + \deg_2(P)$, we have $\overline{\Delta_i}(P) = \overline{\Delta_i}(P)$ by definition of the leading term of a derivation. So $\overline{\Delta_i}(P)$ belongs to I . Since this holds for any element P of I , the derivation $\overline{\Delta_i}$ stabilizes I .

■

The last part of Theorem 1.2 ("Moreover, if this ideal ...") follows from Corollary 2.2.

4 On the parachute for the degree

In this section, we prove Theorem 1.3 using the so-called "parachute" defined in [V07]. Since one wants to majorate $\deg_2(R)$, one may assume that $R \neq 0$. Moreover R cannot be constant so at least one of its partial derivatives $\frac{\partial R}{\partial x_i}$ is nonzero. Without loss of generality we will assume that $\frac{\partial R}{\partial x_n} \neq 0$. Finally recall that $I = (R)$ must be prime so R is irreducible. Let us list our assumptions:

- $\deg_1$ denotes the standard homogeneous degree on $K[x_1, \dots, x_n]$;
- $\Phi = (f_1, \dots, f_n)$ is an automorphism of $K[x_1, \dots, x_n]$;
- $\deg_2$ is the p.w.h. degree defined by $\deg_2(x_i) = d_i = \deg_1(f_i)$, $\forall i = 1, \dots, n$;
- $\bar{P}$ resp. $\tilde{P}$ denotes the leading term of a polynomial with respect to $\deg_1$ resp. $\deg_2$;
- the ideal of relations is principal i.e. $I = (R)$, for some irreducible $R \in K[x_1, \dots, x_n]$ such that $\frac{\partial R}{\partial x_n} \neq 0$.

In our situation the *parachute* of $f_1, \dots, f_n$, denoted ∇ , is simply defined as follows:

$$\nabla := d_1 + \dots + d_n - n.$$

The parachute prevents the degree of a polynomial from "falling too much":

Property 4.1 *For any polynomial P in $K[y_1, \dots, y_m]$, one has the following minoration:*

$$\deg_1(P \circ \Phi) \geq \deg_1\left(\frac{\partial P}{\partial x_n} \circ \Phi\right) + d_n - \nabla \text{ and, inductively,}$$

$$\deg_1(P \circ \Phi) \geq \deg_1\left(\frac{\partial^k P}{\partial x_n^k} \circ \Phi\right) + kd_n - k\nabla, \forall k \geq 0.$$

Proof: It follows from equality (16) applied to $\Phi = (f_1, \dots, f_n)$ that $\deg_1(j(f_1, \dots, f_{n-1}, P \circ \Phi)) = \deg_1(\frac{\partial P}{\partial x_n} \circ \Phi)$. On the other hand, by inequality (15), one has $\deg_1(j(f_1, \dots, f_{n-1}, P \circ \Phi)) \leq \deg_1(f_1) + \dots + \deg_1(f_{n-1}) + \deg_1(P \circ \Phi) - n$ so we have

$$\begin{aligned} \deg_1\left(\frac{\partial P}{\partial x_n} \circ \Phi\right) &\leq d_1 + \dots + d_{n-1} + \deg_1(P \circ \Phi) - n \\ \deg_1\left(\frac{\partial P}{\partial x_n} \circ \Phi\right) - (d_1 + \dots + d_n - n) + d_n &\leq \deg_1(P \circ \Phi) \\ \deg_1\left(\frac{\partial P}{\partial x_n} \circ \Phi\right) + d_n - \nabla &\leq \deg_1(P \circ \Phi) \end{aligned}$$

as required.

■

Lemma 4.2 *Let H be a nonzero polynomial in $K[x_1, \dots, x_n]$ and $k \in \mathbb{N}$ be such that $H \in (R^k) \setminus (R^{k+1})$. Then $\frac{\partial^k H}{\partial x_n^k} \notin (R)$.*

Proof: We proceed by induction on k . For $k = 0$ there is nothing to prove.

Assume this holds for some $k - 1 \geq 0$. One has $H = SR^k$ for some $S \notin R$ and

$$\frac{\partial H}{\partial x_n} = \frac{\partial S}{\partial x_n} R^k + SR^{k-1} \frac{\partial R}{\partial x_n}$$

is clearly in (R^{k-1}) but not in (R^k) because otherwise $S \frac{\partial R}{\partial x_n}$ would be in (R) which is impossible for S by assumption, and for $\frac{\partial R}{\partial x_n}$ for a degree reason so impossible for the product since (R) is prime. The induction assumption concludes. ■

Proposition 4.3 *Let P be a nonzero polynomial in $K[x_1, \dots, x_n]$ and $k \in \mathbb{N}$ be such that $\tilde{P} \in (R^k) \setminus (R^{k+1})$. One has*

- (i) $\deg_1(\frac{\partial^k P}{\partial x^k} \circ \Phi) = \deg_2(\frac{\partial^k P}{\partial x^k}) = \deg_2(P) - kd_n$;
- (ii) $\deg_1(P \circ \Phi) \geq \deg_2(P) - k\nabla$;
- (iii) $\deg_1(P \circ \Phi) \geq k(\deg_2(R) - \nabla)$.

Proof: By Lemma 4.2, $\frac{\partial^k \tilde{P}}{\partial x^k} \notin (R)$. In particular $\frac{\partial^k \tilde{P}}{\partial x^k} \neq 0$ hence, by Lemma 2.6, $\widetilde{\frac{\partial^k \tilde{P}}{\partial x^k}} = \frac{\partial^k \tilde{P}}{\partial x^k} \notin (R) = I$ and $\deg_2(\frac{\partial^k P}{\partial x^k}) = \deg_2(P) - kd_n$ also, by Lemma 2.3, $\deg_1(\frac{\partial^k P}{\partial x^k} \circ \Phi) = \deg_2(\frac{\partial^k P}{\partial x^k}) = \deg_2(P) - kd_n$, thereby proving (i). Assertion (ii) is a direct consequence of (i) and Property 4.1. To prove (iii) observe that $\tilde{P} \in (R^k)$ implies that $k \deg_2(R) \leq \deg_2(\tilde{P}) = \deg_2(P)$. ■

Proof of Theorem 1.3: Let $\Phi^{-1} = (g_1, \dots, g_n)$ be the inverse of Φ . Since $R \neq 0$, it follows from Corollary 2.5 that the automorphism Φ is not affine and neither is its inverse. In particular, there exists an index j such that g_j is not affine. By definition, $x_j = g_j \circ \Phi$ hence $\deg_1(g_j \circ \Phi) = 1$ and, by Corollary 2.4, $\tilde{g}_j \in I = (R)$. It follows that the integer k such that $\tilde{g}_j \in (R^k) \setminus (R^{k+1})$ is positive. Now applying Proposition 4.3 (iii) with $P = g_j$ one has $1 = \deg_1(g_j \circ \Phi) \geq k(\deg_2(R) - \nabla) \geq \deg_2(R) - \nabla$ and consequently $\deg_2(R) \leq 1 + \nabla = d_1 + \dots + d_n - n + 1$. ■

5 On the Jung-van der Kulk Theorem

In this section, we are going to give two elementary proofs of the Jung-van der Kulk Theorem. This theorem states that *every automorphism Φ of K^2 is tame, i.e. Φ is a composition of affine and elementary automorphisms*. Both proofs are based on the computation of the ideal of relations for Φ , and they use induction on the sum $n = \deg_1(f) + \deg_1(g)$, where $\deg_1$ is the standard homogeneous degree on $K[x, y]$ and where $\Phi = (f, g)$.

They proceed as follows. First, if $n = 2$, then Φ is affine and there is nothing to show. So assume that the theorem holds at every order from 2 to n , and let Φ be an automorphism

such that $\deg_1(f) + \deg_1(g) = n + 1$. Up to a permutation of f and g (which corresponds to composing Φ with a linear map), we may assume that $\deg_1(f) \geq \deg_1(g)$. Since Φ is not affine, by Corollary 2.5 the ideal I of relations is distinct from (0) , and its height is either equal to 1 or 2. Note that, if I were of height 2, then it would be equal to (x, y) because I is prime in $K[x, y]$ and weighted homogeneous. But then $\bar{f} = \bar{g} = 0$, which is impossible. So I is prime of height 1, hence it is principal. Write $I = (R)$, where R is irreducible and weighted homogeneous. Assume now that

$$(*) \quad R \text{ is of the form } x - cy^r, \text{ where } c \in K \text{ and } r \in \mathbb{N}^*.$$

Then, this implies that: $\bar{f} - c\bar{g}^r = 0$ and $\deg_1(f - cg^r) < \deg_1(f)$. Consider the map $\Phi' = (f', g')$, where $f' = f - cg^r$ and $g' = g$. As a composition of Φ with an elementary automorphism, it is itself an automorphism of K^2 . Since $\deg_1(f') + \deg_1(g') \leq n$, Φ' is tame by the induction's hypothesis. So Φ is tame and the result follows. Therefore, we only need to prove the assertion $(*)$. In the following subsections, we exhibit two proofs of $(*)$ as an application of our main Theorems.

5.1 First proof of the assertion $(*)$

Let R be an irreducible weighted homogeneous polynomial such that $I = (R)$. Then R is of the form $R(x, y) = \lambda x^s + \mu y^r$, where s, r are coprime. By Theorem 1.2, there exists a nonzero weighted homogeneous locally nilpotent derivation ∂ of $K[x, y]$ such that $\partial(I) \subset I$. In particular, we have $\partial(R) = 0$. One could directly conclude using [ML2] but we prefer to give a self-contained proof. Write ∂ as:

$$\partial = a \frac{\partial}{\partial x} + b \frac{\partial}{\partial y}$$

where a, b are polynomials. Let R^n be the highest power of R that divides both a and b . Since ∂ is locally nilpotent and that $\partial(R) = 0$, $D = \partial/R^n$ is also locally nilpotent. Since $D(R) = 0$ and that $a/R^n, b/R^n$ do not both belong to (R) , D induces a nonzero locally nilpotent derivation D' on the ring:

$$A = \frac{K[x, y]}{(R)} = \frac{K[x, y]}{(\lambda x^s + \mu y^r)} \simeq K[t^r, t^s]$$

Note that, since $\deg_1(f) \geq \deg_1(g)$, we have $r \geq s$. Since s, r are coprime, the integral closure of $K[t^r, t^s]$ is equal to $K[t]$. By Seidenberg's Theorem (see [Sei]), D' extends to a derivation of $K[t]$. By Vasconcelos' Theorem (see [Vas]), this extension is locally nilpotent. So D' is of the form $\theta \partial/\partial t$, where θ belongs to K^* . In particular, $D'(t^s) = s\theta t^{s-1}$ belong to $K[t^r, t^s]$. Since $r \geq s$, this is only possible if $s = 1$, and R is of the form $\lambda x + \mu y^r$.

5.2 Second proof of the assertion $(*)$

Let R be an irreducible weighted homogeneous polynomial such that $I = (R)$. Then R is of the form $R(x, y) = \lambda x^s + \mu y^r$, where s, r are coprime. Set $\deg_1(f) = d_1$ and $\deg_1(g) = d_2$. By assumption, we have $d_2 \leq d_1$. By applying Theorem 1.3 to the polynomial R , we find:

$$sd_2 \leq d_1 + d_2 - 2 < 2d_2$$

Therefore, s is either equal to 0 or 1. But s cannot be equal to 0, otherwise $\bar{g}^r$ would be equal to zero, which is impossible. Therefore, $s = 1$ and $R(x, y) = \lambda x + \mu y^r$.

6 A few surfaces without LND

By Theorem 1.2, in order to show that some polynomials do not appear as generators of relations between the leading terms of an automorphism it is sufficient to show that they are not annihilated by a nontrivial LND. Actually we will show a bit more about those polynomials, namely that the regular functions ring on the surfaces they define admits in turn no nontrivial LND, and use the following

Lemma 6.1 *Let $\deg_1$ be a p.w.h. degree on $K[x_1, \dots, x_n]$. Let R be a $\deg_1$ -homogeneous irreducible element of $K[x_1, \dots, x_n] - K$. Assume there exists a nonzero locally nilpotent derivation D on $K[x_1, \dots, x_n]$ such that $D(R) = 0$. Then the quotient ring $A = K[x_1, \dots, x_n]/(R)$ admits a nonzero weighted homogeneous locally nilpotent derivation.*

Proof: Since D is locally nilpotent, its leading term $\overline{D}$ is nonzero and locally nilpotent (see [vdE]). Since R is weighted homogeneous and that $D(R) = 0$, we have $\overline{D}(R) = 0$. So we may assume that D is weighted homogeneous. Write D as:

$$D = \sum_{i=1}^n a_i \frac{\partial}{\partial x_i}.$$

Let R^k be the maximal common power of R dividing $a_1, \dots, a_n$. Since $D(R) = 0$, the derivation $\partial = D/R^k$ is a locally nilpotent derivation on $K[x_1, \dots, x_n]$ which induces a weighted homogeneous locally nilpotent derivation ∂' on the quotient ring A . Moreover, since not all a_i/R^k are divisible by R , ∂' is nonzero. ■

Using some results and technics of [KZ00] we prove

Proposition 6.2 *The quotient algebra $K[x_1, x_2, x_3]/(R)$ admit no nonzero LND when R is one of the following polynomials:*

$x_3^2 + ax_1^4 + bx_2^3$	where	$ab \neq 0$
$x_3^2 + ax_1^5 + bx_2^3$	where	$ab \neq 0$
$x_3^2 + (a_1x_1 + b_1x_2)(a_2x_1 + b_2x_2)(a_3x_1 + b_3x_2)$	where	$a_ib_j - a_jb_i \neq 0, \forall i \neq j$
$x_3^2 + (a_1x_1^{e_1} + b_1x_2)(a_2x_1^{e_1} + b_2x_2)x_1$	where	$a_1b_2 - a_2b_1 \neq 0, e_1 \geq 0$
$x_3^2 + (ax_1^3 + bx_2^2)x_2$	where	$ab \neq 0$

Proof: Up to multiplication of x_1 and x_2 by appropriate scalars and renaming the indeterminates, our first two polynomials define the Platonic surfaces $S_{2,3,4}$ and $S_{2,3,5}$ where $S_{k,l,m} = \{x^k + y^l + z^m = 0\} \subset K^3$ as in the proof of Lemma 4 in [KZ00]. Even if, in the cited article, K is the complex field, the proofs that $S_{2,3,4}$ and $S_{2,3,5}$ admit no non-trivial LND still works word for word for a field as general as ours. Our third polynomial defines a surface isomorphic to $S_{2,3,3}$: in a word one chooses a linear change of the coordinates x_1, x_2 corresponding to the projective map taking (a_i, b_i) to $(1, \lambda^i)$ for $i = 1, 2, 3$ where λ is a primitive cubic root of the unity and one rescales x_3 . Again by [KZ00], $S_{2,3,3}$ admit no non-trivial LND. Now we assume that R is one of the two last polynomials with weights $(\deg_1(x_1), \deg_1(x_2), \deg_1(x_3)) = (d_1, d_2, d_3) = (2, 2e_1, 2e_1 + 1)$ resp. $(4, 6, 9)$, and that the quotient ring $A = K[x_1, x_2, x_3]/(R)$ admits a nonzero LND ∂ . Up to taking the leading term of ∂ as in [KZ00] (see also the beginning of our section 3) one may assume that this derivation is weighted homogeneous.

First, we are going to prove that $\ker \partial$ contains an irreducible weighted homogeneous polynomial $P(x_1, x_2)$. Since A has transcendence degree 2 over K , there exists a nonconstant element f of A such that $\partial(f) = 0$. Since A is graded and that ∂ is weighted homogeneous, we may assume that f is also weighted homogeneous. Consider R as a polynomial of degree 2 in x_3 . After division by R , we may assume that f is of the form:

$$f = f_1(x_1, x_2)x_3 + f_0(x_1, x_2)$$

where f_1, f_0 are polynomials. Since $\deg_1(x_i)$ is even for $i = 1, 2$, $\deg_1(f_1)$ and $\deg_1(f_2)$ are even. Since f is weighted homogeneous and $\deg_2(x_3)$ is odd, then either $f = f_1(x_1, x_2)x_3$ or $f = f_0(x_1, x_2)$. In all cases, since x_3^2 is a polynomial in x_1, x_2 , f^2 belongs to $\ker \partial \cap K[x_1, x_2]$. So $\ker \partial$ contains a nonconstant weighted homogeneous element P of $K[x_1, x_2]$. Since ∂ is locally nilpotent, $\ker \partial$ is factorially closed (well-known fact, see e.g. [ML1]) and every irreducible factor of P in $K[x_1, x_2]$ lies in $\ker \partial$. So we may assume that P is irreducible.

Second, note that, since P is weighted homogeneous, it is either equal (up to a scalar multiple) to x_1 , or x_2 , or $cx_1^r + x_2^s$, where $c \in k$. Now let us distinguish between the two cases.

If $R = x_3^2 + (a_1x_1^{e_1} + b_1x_2)(a_2x_1^{e_1} + b_2x_2)x_1$ with the weights $(2, 2e_1, 2e_1 + 1)$ and $P = cx_1^r + x_2^s$ then $r = e_1$ and $s = 1$ so that it is weighted homogeneous. The change of coordinates $(x_1, x_2, x_3) \mapsto (x_1, x_2 - cx_1^r, x_3)$ doesn't affect the form of R , so we may assume $c = 0$. So we are left with the two cases $x_1 \in \ker \partial$ or $x_2 \in \ker \partial$.

Assume that $\partial(x_1) = 0$, and let L be the algebraic closure of the field $K(x_1)$. Define a new derivation D on the ring $B = L \otimes_{K[x_1]} A$ as:

$$D : L \otimes_{K[x_1]} A \longrightarrow L \otimes_{K[x_1]} A, \quad a \otimes b \longmapsto a \otimes \partial(b).$$

Since ∂ is nonzero locally nilpotent, D is also nonzero locally nilpotent. Moreover, B is isomorphic to the ring $L[x_2, x_3]/(x_3^2 + x_2^2 - 1)$. Since $\text{Spec}(B)$ is isomorphic to L^* , D is equal to zero, which is impossible. So $\partial(x_1) \neq 0$ and $\partial(x_2) = 0$. Then, following the same construction, we obtain a nonzero locally nilpotent derivation D on the coordinate ring of the curve $x_3^2 + (a_1x_1^{e_1} + b_1x_2)(a_2x_1^{e_1} + b_2x_2)x_1 = 0$ over L , where L is the algebraic closure of the field $K(x_2)$. Since this curve is smooth and hyperelliptic, its geometric genus is equal to $e_1 - 1$. In particular, this curve cannot be isomorphic to L unless $e_1 = 1$. But in this case we get L^* ; therefore D is equal to 0, hence a contradiction.

If $R = x_3^2 + (ax_1^3 + bx_2^2)x_2$ with the weights $(4, 6, 9)$ then the case $P = cx_1^r + x_2^s$ is excluded since (r, s) would have to be $(3, 2)$ which is impossible by [ML2]. It follows that x_1 or x_2 is in $\text{Ker} \partial$ and we get a contradiction as in the previous case.

■

7 A special surface without LND

In this section, we are going to establish the following result:

Proposition 7.1 *Let e_1 be in $3 + 2\mathbb{N}$, $a, b \in K^*$ and $R = x_3^2 + (ax_1^{e_1} + bx_2^2)x_1$. Then the quotient algebra $K[x_1, x_2, x_3]/(R)$ admit no nonzero LND.*

Up to replacing each x_i with $\lambda_i x_i$ for some suitable constants λ_i , we may assume that $a = 1$ and $b = -1$. For any element e_1 of $3 + 2\mathbb{N}$, denote by S the surface of K^3 given by the equation $R = 0$. Consider the Danielewski surface S' of K^3 , given by the equation:

$$\alpha^{2e_1} - 4\beta\gamma = 0$$

together with the involution σ of S' defined by the formula $\sigma(\alpha, \beta, \gamma) = (-\alpha, \gamma, \beta)$. Denote by σ^* the involution of $\mathcal{O}(S')$ defined by $\sigma^*(f) = f \circ \sigma$. It is easy to check that σ has $(0, 0, 0)$ as a unique fixpoint, and that the map:

$$F : S' \longrightarrow S, \quad (\alpha, \beta, \gamma) \longmapsto (\alpha^2, \beta + \gamma, \alpha(\beta - \gamma))$$

is a well-defined morphism, i.e. $F(S') \subseteq S$. Since F is dominant, it induces an injective algebra morphism from $\mathcal{O}(S)$ to $\mathcal{O}(S')$. We can therefore consider $\mathcal{O}(S)$ as a subring of $\mathcal{O}(S')$. Note that S' is weighted homogeneous for the weights $1, e_1, e_1$. Moreover, the coordinate functions of F are all weighted homogeneous of degree $2, e_1, e_1 + 1$. We begin with a few lemmas.

Lemma 7.2 *Both surfaces S and S' are normal.*

Proof: Since S and S' are hypersurfaces of K^3 , it suffices to show that they are both nonsingular in codimension 1 by Serre's criterion. Since $(0, 0, 0)$ is their only singular point, they are normal. ■

Lemma 7.3 *The morphism F is the quotient map of the $\mathbb{Z}/2\mathbb{Z}$ -action on S' defined by the involution σ . In particular, $S = S' / (\mathbb{Z}/2\mathbb{Z})$.*

Proof: By means of the morphism F , we can identify $\mathcal{O}(S)$ with a subring of $\mathcal{O}(S')$. Via this identification, we only need to show that $\mathcal{O}(S)$ is the ring of invariants of σ^* . We proceed in several steps. First, we show that F is a finite morphism. Note that α^2 and $\beta + \gamma$ belong to $\mathcal{O}(S)$, so that α and $\beta + \gamma$ are integral over $\mathcal{O}(S)$. Since $\alpha^{2e_1} - 4\beta\gamma = 0$, we obtain the relation:

$$(\beta - \gamma)^2 = -\alpha^{2e_1} + (\beta + \gamma)^2.$$

In particular, $\beta - \gamma$ is integral over $\mathcal{O}(S)$. Therefore, α , β and γ are integral over $\mathcal{O}(S)$, hence the morphism F is finite. Second, denote by $K(S)$ (resp. $K(S')$) the field of rational functions of S (resp. S'). We are going to prove that $K(S)$ is the field $K(S')^{\sigma^*}$ of invariants of σ^* . It is easy to check that the functions α^2 , $\beta + \gamma$ and $\alpha(\beta - \gamma)$ are σ^* -invariant. In particular, $K(S)$ is contained in $K(S')^{\sigma^*}$. Since α^2 belongs to $K(S)$ and that $K(S') = K(S)[\alpha]$, $K(S')/K(S)$ is an extension of degree 2. In particular, we find that $K(S')^{\sigma^*} = K(S)$. Now, we can prove that $\mathcal{O}(S')^{\sigma^*} = \mathcal{O}(S)$. Since the coordinate functions of F are σ^* -invariant, $\mathcal{O}(S)$ is contained in $\mathcal{O}(S')^{\sigma^*}$. Conversely, let f be any σ^* -invariant regular function on S' . Since $K(S')^{\sigma^*} = K(S)$, f belongs to $K(S)$. Since F is finite, f is integral over $\mathcal{O}(S)$. But S is normal, so $\mathcal{O}(S)$ is integrally closed and f belongs to $\mathcal{O}(S)$. ■

Lemma 7.4 *Let D be any derivation on $\mathcal{O}(S)$. Then there exists a unique derivation D' on $\mathcal{O}(S')$, which commutes with σ^* and such that D' coincides with D on $\mathcal{O}(S)$. Moreover, if D is weighted homogeneous on S , then D' is also weighted homogeneous on S' .*

Proof: By assumption, D is a K -derivation on $\mathcal{O}(S)$, hence it extends uniquely into a K -derivation of $K(S)$. Since $K(S')/K(S)$ is a finite extension, D extends uniquely into a K -derivation D' of $K(S')$ by some elementary results of Differential Galois Theory (see [Ko]). Consider the map $D_0 = \sigma^* \circ D' \circ \sigma^*$ on $K(S')$. By construction, D_0 is a K -linear map on $K(S')$. Since σ^* is an involution, we have for any elements f and g of $K(S')$:

$$\begin{aligned} D_0(fg) &= \sigma^*(D(\sigma^*(f)\sigma^*(g))) \\ &= \sigma^*(\sigma^*(f)D(\sigma^*(g)) + \sigma^*(\sigma^*(g)D(\sigma^*(f)))) \\ &= fD_0(g) + gD_0(f). \end{aligned}$$

Therefore, D_0 satisfies the Leibniz rule, hence it is a K -derivation of $K(S')$. Since σ^* is the identity on $K(S)$, D coincides with D_0 on $K(S)$. By uniqueness of the extension of a K -derivation in an algebraic extension, we find that $D = D_0$, hence $D' \circ \sigma^* = \sigma^* \circ D'$. So there only remains to prove that D' preserves the ring $\mathcal{O}(S')$.

Let p be any point of S' distinct from $(0,0,0)$. Then the point $q = F(p)$ is distinct from $(0,0,0)$. In particular, p (resp. q) is a smooth point of S' (resp. S). Since F is a quotient map of a finite group of order 2, its singular points correspond to its fixpoints. Since $(0,0,0)$ is the only fixpoint of σ^* , F is nonsingular at p . In particular, the pull-back morphism F^* defines an isomorphism between the completions of the local rings $\mathcal{O}_{S',p}$ and $\mathcal{O}_{S,q}$ with respect to their maximal ideals. Via this isomorphism, D determines a unique derivation D'' on the completion of $\mathcal{O}_{S',p}$. By uniqueness of the extension of D , we deduce that $D' = D''$. In particular, D' is regular at the point p . Since this holds for any point $p \neq (0,0,0)$, D' is regular on $S' - \{(0,0,0)\}$. Thus, for any regular function f on S' , $D'(f)$ is regular on $S' - \{(0,0,0)\}$. But since S' is normal, $D'(f)$ is regular on all of S' . In particular, D' preserves the ring $\mathcal{O}(S')$.

Finally, assume that D is weighted homogeneous on S . By definition, this means there exists a constant r such that D maps every weighted homogeneous element of degree n to a weighted homogeneous element of degree $n+r$. Since α^2 is a weighted homogeneous element of $\mathcal{O}(S)$ of degree 2, $D'(\alpha) = D(\alpha^2)/2\alpha$ is weighted homogeneous of degree $r+1 = r + \deg(\alpha)$. Since $\beta + \gamma$ is a weighted homogeneous element of $\mathcal{O}(S)$ of degree e_1 , $D'(\beta + \gamma) = D(\beta + \gamma)$ is weighted homogeneous of degree $r + e_1 = r + \deg(\beta + \gamma)$. Similarly, one can check that $D'(\beta - \gamma)$ is weighted homogeneous of degree $e_1 + r = r + \deg(\beta - \gamma)$. Since α , $\beta + \gamma$ and $\beta - \gamma$ span the graded ring $\mathcal{O}(S')$, D' is a weighted homogeneous K -derivation of degree r . ■

Proof of Proposition 7.1: Assume that the polynomial $R = x_3^2 + (x_1^{e_1} - x_2^2)x_1$ is annihilated by a nonzero locally nilpotent derivation. By Lemma 6.1, the surface S admits a nonzero weighted homogeneous locally nilpotent derivation D . By Lemma 7.4, D extends uniquely into a weighted homogeneous K -derivation D' of S' , which commutes with the involution σ^* . Since $\mathcal{O}(S')$ is integral over $\mathcal{O}(S)$ by Lemma 7.3 and that D is locally nilpotent, D' is also locally nilpotent by Vasconcelos' Theorem (see [Vas]). Set $\varphi(x) = x^{2e_1}/4$. For any polynomial f of $K[x]$, define as in [Dai] the automorphism Δ_f of S' by the formula:

$$\Delta_f(\alpha, \beta, \gamma) = \left(\alpha + \beta f(\beta), \beta, \frac{\varphi(\alpha + \beta f(\beta))}{\beta} \right).$$

Let δ be the automorphism of S' given by $\delta(\alpha, \beta, \gamma) = (\alpha, \gamma, \beta)$. Denote by G the group spanned by δ and all the Δ_f , where f runs through $K[x]$. Then G acts transitively on the kernels of

nonzero locally nilpotent derivations on S' (see [Dai]). Since $K[\beta]$ is the kernel of a locally nilpotent derivation on S' , this means that $\ker D'$ must be of the form $K[g]$, where g is the second coordinate function of an element of G . Note that every generator of G fixes the origin $(0, 0, 0)$, and that its linear part at $(0, 0, 0)$ is either of the form $(\alpha, \beta, \gamma) \mapsto (\alpha + a\beta + b\gamma, \beta, \gamma)$ or $(\alpha, \beta, \gamma) \mapsto (\alpha + a\beta + b\gamma, \gamma, \beta)$. So every element of G fixes the point $(0, 0, 0)$, and its linear part at $(0, 0, 0)$ is either of the form $(\alpha, \beta, \gamma) \mapsto (\alpha + a\beta + b\gamma, \beta, \gamma)$ or $(\alpha, \beta, \gamma) \mapsto (\alpha + a\beta + b\gamma, \gamma, \beta)$. Therefore, g must be either of the form $\beta + h$ or $\gamma + h$, where h is a polynomial with no constant nor linear terms in α, β, γ . Since D' is weighted homogeneous and that $g(0, 0, 0) = 0$, g must be also weighted homogeneous. So g is either of the form $\beta + \lambda\alpha^{e_1}$ or $\gamma + \lambda\alpha^{e_1}$, where λ belongs to K . Now, since D' commutes with σ^* , $\ker D'$ must be stable by the action of σ^* . In particular, g is σ^* -equivariant. But this is impossible since σ^* permutes β and γ and that $\sigma^*(\alpha) = -\alpha$. This ends the proof of the proposition. ■

Remark 7.5 One has $S' \simeq K^2 // (\mathbb{Z}/2e_1\mathbb{Z})$ where $\mathbb{Z}/2e_1\mathbb{Z}$ acts on K^2 through $(x, y) \mapsto (-\xi x, -\xi^{-1}y)$, ξ being a primitive e_1 -th root of the unity. In turn, the action on K^2 induced by $(x, y) \mapsto (\xi x, \xi^{-1}y)$ and $(x, y) \mapsto (-y, x)$ gives $S \simeq S' // (\mathbb{Z}/2\mathbb{Z}) = K^2 // G$ where G is a semi-direct product $\mathbb{Z}/e_1\mathbb{Z} \ltimes \mathbb{Z}/4\mathbb{Z}$. In case $K = \mathbb{C}$ the fact that S does not admit a non-trivial LND ($\mathbb{C}_+$ -action) follows from Theorem 2.13 in [FZ]: G is not cyclic and the quotient $\mathbb{C}^2 // G$ is surely not isomorphic to a quotient of $\mathbb{C}^2$ by a cyclic group (the group by which one takes the quotient reflects in the fundamental group of the smooth locus of the surface).

8 Proof of the third Theorem

Assume that the ideal I of relations is nonzero principal, i.e. $I = (R)$ with $R \neq 0$. By Theorem 1.2, Lemma 6.1, Proposition 6.2 and 7.1 R is proportional to none of the polynomials in the following "forbidden" list:

$$\begin{array}{ll}
 x_3^2 + ax_1^4 + bx_2^3 & \text{where } ab \neq 0 \\
 x_3^2 + ax_1^5 + bx_2^3 & \text{where } ab \neq 0 \\
 x_3^2 + (a_1x_1^{e_1} + b_1x_2)(a_2x_1^{e_1} + b_2x_2)x_1 & \text{where } a_1b_2 - a_2b_1 \neq 0, e_1 \geq 1 \\
 x_3^2 + (a_1x_1 + b_1x_2)(a_2x_1 + b_2x_2)(a_3x_1 + b_3x_2) & \text{where } a_ib_j - a_jb_i \neq 0, \forall i \neq j \\
 x_3^2 + (ax_1^3 + bx_2^2)x_2 & \text{where } ab \neq 0 \\
 x_3^2 + (ax_1^{e_1} + bx_2^2)x_1 & \text{where } ab \neq 0, e_1 \in 3 + 2\mathbb{N}.
 \end{array}$$

Being homogeneous, the polynomial R can be written as:

$$R = \sum_{\alpha_1 d_1 + \alpha_2 d_2 + \alpha_3 d_3 = \deg_2(R)} R_{\alpha_1, \alpha_2, \alpha_3} x_1^{\alpha_1} x_2^{\alpha_2} x_3^{\alpha_3}.$$

Denote by $\text{Supp}(R)$ the set $\{\alpha = (\alpha_1, \alpha_2, \alpha_3) | R_\alpha \neq 0\}$. Then Theorem 1.3 implies:

$$(17) \quad \forall \alpha \in \text{Supp}(R), \quad \alpha_1 d_1 + \alpha_2 d_2 + \alpha_3 d_3 \leq d_1 + d_2 + d_3 - 2$$

Assume that $d_1 \leq d_2 \leq d_3$. Since $d_1 + d_2 + d_3 - 2 < 3d_3$, we find that $\alpha_3 \leq 2$ for any element α of $\text{Supp}(R)$. In particular, R is a polynomial of degree ≤ 2 in the variable x_3 . Set $d = \gcd(d_1, d_2)$ and write $d_1 = de_2$, $d_2 = de_1$. Note that $\gcd(e_1, e_2) = 1$ and $e_2 \leq e_1$, with equality if and only if $e_1 = e_2 = 1$.

8.1 The case when $\deg_{x_3}(R) = 0$

Then, R is an irreducible weighted homogeneous polynomial in the variables x_1, x_2 and for the weights d_1, d_2 . Since K is algebraically closed, R is either of the form $ax_1^{e_1} + bx_2^{e_2}$ or cx_i , where $i \in \{1, 2\}$ and $a, b, c \neq 0$. Since neither $\bar{f}_1$ nor $\bar{f}_2$ is zero, R has type (3).

8.2 The case when $\deg_{x_3}(R) = 1$

Let α be any point of $\text{Supp}(R)$ of the form $\alpha = (\alpha_1, \alpha_2, 1)$. By inequality (17), we have:

$$\begin{aligned} \alpha_1 d_1 + \alpha_2 d_2 + d_3 &\leq d_1 + d_2 + d_3 - 2 \Rightarrow \alpha_1 d_1 + \alpha_2 d_2 \leq d_1 + d_2 - 2 \\ &\Rightarrow \alpha_2 \leq 1 \text{ and } (\alpha_2 = 1 \Rightarrow \alpha_1 = 0) . \end{aligned}$$

If $(0, 1, 1)$ does not belong to $\text{Supp}(R)$, then α_2 is always zero and R has type (5) with $x'_3 = x_3$. Now assume that $(0, 1, 1)$ belongs to $\text{Supp}(R)$. Then, up to a multiplication by a constant, R is of the form:

$$R = (x_2 + ax_1^{e_1})x_3 + P(x_1, x_2)$$

where $a \neq 0$ only if $e_2 = 1$, and where $P(x_1, x_2)$ is weighted homogeneous. Write P as $P(x_1, x_2) = P_0(x_1) + (x_2 + ax_1^{e_1})Q(x_1, x_2 + ax_1^{e_1})$. Taking $x'_3 = x_3 + Q(x_1, x_2 + ax_1^{e_1})$, we get a polynomial of type (4).

8.3 The case when $\deg_{x_3}(R) = 2$

Let α be any element of $\text{Supp}(R)$ of the form $(\alpha_1, \alpha_2, 2)$. Assume that either α_1 or α_2 is nonzero. Then, by inequality (17), we would find:

$$d_1 + 2d_3 \leq \alpha_1 d_1 + \alpha_2 d_2 + 2d_3 \leq d_1 + d_2 + d_3 - 2 \Rightarrow d_3 \leq d_2 - 2$$

which is impossible. So, if $\alpha = (\alpha_1, \alpha_2, 2)$, then α is equal to $(0, 0, 2)$ and

$$\begin{aligned} 2d_3 &\leq d_1 + d_2 + d_3 - 2 \Rightarrow \\ (18) \quad d_3 &\leq d_1 + d_2 - 2 . \end{aligned}$$

This means in particular that, up to a scalar multiplication, R is of the form:

$$R = x_3^2 + x_3 P(x_1, x_2) + Q(x_1, x_2)$$

where P and Q are weighted homogeneous. Taking $x'_3 = x_3 + P(x_1, x_2)/2$, we have:

$$\begin{aligned} R &= x_3'^2 + \text{a weighted homogeneous polynomial in } x_1, x_2 \\ &= x_3'^2 + c \prod_{i=1}^k (a_i x_1^{e_1} + b_i x_2^{e_2}) x_1^{r_1} x_2^{r_2} \end{aligned}$$

where $r_l < e_l$ for $l = 1, 2$ (c is assumed to be one as soon as $k \geq 1$). Since R is weighted homogeneous, we obtain:

$$\begin{aligned} 2d_3 &= ke_k d_k + r_1 d_1 + r_2 d_2 \text{ for } k = 1, 2 \\ &= kde_1 e_2 + r_1 d_1 + r_2 d_2 . \end{aligned}$$

Inequality (18) yields $d_3 \leq de_1 + de_2 - 2$ and since $d_2 \leq d_3$ one gets:

$$2d_2 = 2de_1 \leq 2d_3 = kde_1e_2 + r_1de_2 + r_2de_1 \leq 2de_1 + 2de_2 - 4.$$

After division by de_1e_2 , this yields:

$$(19) \quad 2/e_2 \leq k + r_1/e_1 + r_2/e_2 < 2/e_2 + 2/e_1.$$

In particular, we have $k < 4$. We are going to describe the different possible polynomials R depending on the value of k .

1st case: $k = 0$ i.e. $R = x_3'^2 + cx_1^{r_1}x_2^{r_2}$.

Inequalities (19) becomes $2/e_2 \leq r_1/e_1 + r_2/e_2 < 2/e_2 + 2/e_1$ and we deduce that either $r_1 \leq 1$ or $r_2 \leq 1$.

If $r_1 = 0$, then we have:

$$2/e_2 \leq r_2/e_2 < 2/e_2 + 2/e_1.$$

After multiplication by e_2 , we find $2 \leq r_2 < 2 + 2e_2/e_1$. Since $e_2/e_1 \leq 1$, we obtain that either $r_2 = 2$ or $r_2 = 3$. Since R is irreducible, the case $r_2 = 2$ is impossible and we have:

$$R = x_3'^2 + cx_2^3.$$

So R corresponds to a polynomial of type (6) in the list.

If $r_1 = 1$, then we have:

$$\begin{aligned} 2/e_2 \leq 1/e_1 + r_2/e_2 < 2/e_2 + 2/e_1 &\Rightarrow 2e_1/e_2 \leq 1 + r_2e_1/e_2 < 2 + 2e_1/e_2 \\ &\Rightarrow 2e_1/e_2 - 1 \leq r_2e_1/e_2 < 2e_1/e_2 + 1 \\ &\Rightarrow -1 \leq (r_2 - 2)e_1/e_2 < 1. \end{aligned}$$

Since $e_1/e_2 \geq 1$, r_2 is either equal to 1 or 2. Therefore R is either of type (9) or (7).

Now, if $r_1 \geq 2$, then r_2 is either equal to 0 or 1. If $r_2 = 0$, then, by irreducibility, $r_1 \in 3 + 2\mathbb{N}$ and R is of type (8). If $r_2 = 1$, then R is of type (9).

2nd case: $k = 1$ i.e. $R = x_3'^2 + (ax_1^{e_1} + bx_2^{e_2})x_1^{r_1}x_2^{r_2}$.

Inequalities (19) become $2/e_2 \leq 1 + r_1/e_1 + r_2/e_2 < 2/e_2 + 2/e_1$ and it follows that either $r_1 \leq 1$ or $r_2 \leq 1$.

If $r_1 = 0$, then we have:

$$2/e_2 \leq 1 + r_2/e_2 < 2/e_2 + 2/e_1.$$

After multiplication by e_2 , we find $2 \leq e_2 + r_2 < 2 + 2e_2/e_1$. Since $e_2/e_1 \leq 1$, $e_2 + r_2$ is either equal to 2 or 3. Assume that $e_2 + r_2 = 2$. Since $r_2 < e_2$, we have $r_2 = 0$ and $e_2 = 2$. Therefore, R is of type (10). Assume now that $e_2 + r_2 = 3$. Then we find that $1 < 2e_2/e_1$, hence $e_1 < 2e_2$. Recalling that $e_2 \leq e_1$ with equality only if $e_1 = e_2 = 1$, we are left with the following three cases:

$$\begin{cases} k = 1, r_1 = 0, r_2 = 1, e_2 = 2, e_1 = 3 \\ k = 1, r_1 = 0, r_2 = 0, e_2 = 3, e_1 = 4 \\ k = 1, r_1 = 0, r_2 = 0, e_2 = 3, e_1 = 5 \end{cases}.$$

Therefore, R is equal to one of the following polynomials:

$$\begin{cases} x_3'^2 + (ax_1^3 + bx_2^2)x_2 \\ x_3'^2 + ax_1^4 + bx_2^3 \\ x_3'^2 + ax_1^5 + bx_2^3. \end{cases}$$

They all belong to the forbidden list at the beginning of this section unless $ab = 0$. Since $x_3'^2 + ax_1^4$ is reducible, R has either type (9), (6) or (8).

If now $r_1 = 1$, then we have:

$$2/e_2 \leq 1 + 1/e_1 + r_2/e_2 < 2/e_2 + 2/e_1.$$

After multiplication by e_2 , we obtain:

$$2 \leq e_2 + r_2 + \frac{e_2}{e_1} < 2 + 2\frac{e_2}{e_1} \Rightarrow 2 - \frac{e_2}{e_1} \leq e_2 + r_2 < 2 + \frac{e_2}{e_1}.$$

Since $e_2/e_1 \leq 1$, we find that $e_2 + r_2$ is either equal to 1 or 2. Assume that $e_2 + r_2 = 1$. Then $2 - e_2/e_1 = 1$ i.e. $e_1 = e_2 = 1$ and, since $r_2 < e_2$, we obtain that $r_2 = 0$. Therefore, R is of the form $x_3'^2 + (ax_1 + bx_2)x_1$, i.e. it is of type (11). Assume now that $e_2 + r_2 = 2$. Since $r_2 < e_2$, we find that $r_2 = 0$ and $e_2 = 2$. Therefore, recalling that $k = 1$ and $e_1 > r_1 = 1$, R is of the form $R = x_3'^2 + (ax_1^{e_1} + bx_2^2)x_1$ with $1 = \gcd(e_1, e_2) = \gcd(e_1, 2) \Rightarrow e_1 \in 3 + 2\mathbb{N}$. So R belongs to the forbidden list unless $ab = 0$. If $a = 0$ and $b \neq 0$, then R is of type (7). If $a \neq 0$ and $b = 0$, then R is reducible.

Finally assume $r_1 \geq 2$. The right hand side inequality in (19) yields

$$\begin{aligned} 1 + 2/e_1 + r_2/e_2 &< 2/e_2 + 2/e_1 \Rightarrow 1 + r_2/e_2 < 2/e_2 \\ &\Rightarrow e_2 + r_2 < 2. \end{aligned}$$

Since $r_2 < e_2$, we get that $k = 1$, $r_1 \geq 2$, $r_2 = 0$ and $e_2 = 1$. Therefore, R is of the form $x_3'^2 + (ax_1^{e_1} + bx_2)x_1^{r_1}$, i.e. it is of type (11).

3rd case: $k = 2$ i.e. $R = x_3'^2 + (a_1x_1^{e_1} + b_1x_2^{e_2})(a_2x_1^{e_1} + b_2x_2^{e_2})x_1^{r_1}x_2^{r_2}$.

We start from (19):

$$\begin{aligned} 2/e_2 \leq 2 + r_1/e_1 + r_2/e_2 &< 2/e_2 + 2/e_1 \Rightarrow 2 < 2/e_2 + 2/e_1 \leq 4/e_2 \\ &\Rightarrow e_2 < 2 \end{aligned}$$

so we get $e_2 = 1$, $r_2 = 0$ and, using again the left hand side of the implication: $2 \leq 2 + r_1/e_1 < 2 + 2/e_1$ which simply means $r_1 < 2$. Therefore, R is equal to one of the following polynomials:

$$\begin{cases} x_3'^2 + (a_1x_1^{e_1} + b_1x_2)(a_2x_1^{e_1} + b_2x_2) \\ x_3'^2 + (a_1x_1^{e_1} + b_1x_2)(a_2x_1^{e_1} + b_2x_2)x_1 \end{cases}.$$

The first one is of type (12) when not reducible.

The second one is in the forbidden list unless $a_1b_2 - a_2b_1 = 0$, which gives a polynomial of the form $x_3'^2 + (ax_1^{e_1} + bx_2)^2x_1$, i.e. of type (14).

4th case: $k = 3$ i.e. $R = x_3'^2 + (a_1x_1^{e_1} + b_1x_2^{e_2})(a_2x_1^{e_1} + b_2x_2^{e_2})(a_3x_1^{e_1} + b_3x_2^{e_2})x_1^{r_1}x_2^{r_2}$.

Again (19) yields

$$2/e_2 \leq 3 + r_1/e_1 + r_2/e_2 < 2/e_2 + 2/e_1 \Rightarrow 3 < 2/e_2 + 2/e_1$$

which is possible only if $e_1 = e_2 = 1$. Consequently $r_1 = r_2 = 0$ and R is of the form:

$$R = x_3'^2 + (a_1x_1 + b_1x_2)(a_2x_1 + b_2x_2)(a_3x_1 + b_3x_2).$$

This is again in the forbidden list unless one of the determinants $a_ib_j - a_jb_i$ is zero. In this case, R is of type (13), and the classification is complete.

9 Proof of the Corollary

Proof of Corollary 1.5: By Theorem 1.4 we have to prove that every one of the 14 polynomials in the list given inside this theorem is tamely equivalent to 0, x_3 , $x_1^r + x_2^s$ or $x_1^k x_3 + P(x_1, x_2)$. Let us introduce the notation $Q \equiv R$ when Q and R are tamely equivalent and we call R_i , $\forall i = 1, \dots, 14$, the polynomial such the relation (i) in the list is equal to $R_i(x_1, x_2, x_3')$. Applying the elementary automorphism $(x_1, x_2, x_3 - h(x_1, x_2))$ we get $R_i(x_1, x_2, x_3') \equiv R_i(x_1, x_2, x_3)$ so we may forget the "prime" in x_3' .

We have $R_1 = 0$ and $R_2 = x_3$ so there is nothing to do. One has $R_3 \circ (x_1, dx_2, x_3) = x_1^{e_1} + x_2^{e_2}$ where $d^s = 1/c$ so $R_3 \equiv x_1^r + x_2^s$ with $(r, s) = (e_1, e_2)$. One has $R_4 \circ (x_1, x_2 + ax_1^{e_1}, x_3) = R_3 \equiv x_1^r + x_2^s$ so $R_4 \equiv x_1^r + x_2^s$. The relation R_5 is even equal to $x_1^k x_3 + P(x_1, x_2)$. By a transposition (x_3, x_2, x_1) we get $R_6 \equiv R_3 \equiv x_1^r + x_2^s$ with $(r, s) = (2, 3)$. Similarly, using if needed some permutation and some tame automorphism of the form $(x_1, \frac{1}{b}(x_2 - ax_1^{e_1}))$ one gets that R_7 , R_9 , R_{11} and R_{14} are tamely equivalent to $x_1^k x_3 + P(x_1, x_2)$ and R_8 to $x_1^r + x_2^s$. As for R_{10} , let $d^2 = -b$ and compose $R_{10} \circ (x_1, \frac{x_2 - x_3}{2d}, \frac{x_2 + x_3}{2}) = ax_1^{e_1} + x_3x_2 \equiv x_1^k x_3 + P(x_1, x_2)$. In R_{12} , if $b_1b_2 = 0$ then clearly $R_{12} \equiv R_{11}$ or directly R_9 , that is, equivalent to $x_1^k x_3 + P(x_1, x_2)$. If $b_1b_2 \neq 0$ then one checks that $R_{12} \circ (x_1, x_2 - \frac{a_1b_2 + b_1a_2}{2b_1b_2}x_1^{e_1}, x_3)$ looks like R_{10} but with an even e_1 , which does not prevent us from showing, as before, that $R_{12} \equiv x_1^k x_3 + P(x_1, x_2)$. Finally, using an appropriate linear automorphism shows that $R_{13} \equiv R_7 \equiv x_1^k x_3 + P(x_1, x_2)$ unless $a_1b_2 - a_2b_1 = 0$, in which case $R_{13} \equiv R_3$. ■

Remark that the four polynomials in Corollary 1.5 do belong to kernels of nontrivial LNDs of $K[x_1, x_2, x_3]$: 0 and x_3 are in the kernel of $\frac{\partial}{\partial x_1}$, $x_1^r + x_2^s$ in the one of $\frac{\partial}{\partial x_3}$ and $x_1^k x_3 + P(x_1, x_2)$ is in the kernel of $x_1^k \frac{\partial}{\partial x_2} - \frac{\partial P}{\partial x_2} \frac{\partial}{\partial x_3}$.

References

- [Dai] D. Daigle, *On locally nilpotent derivations of $k[X_1, X_2, Y]/(\varphi(Y) - X_1X_2)$* , J. Pure Appl. Algebra **181** (2003), n° 2-3, 181-208.
- [vdE] A. van den Essen, *Polynomial automorphisms and the Jacobian conjecture*, Progress in Mathematics 190, Birkhäuser Verlag, Basel, 2000.
- [FZ] H. Flenner and M. Zaidenberg, *Rational curves and rational singularities*, Math. Z. 244, No. 3, 549-575 (2003).
- [Jun] H. Jung, *Über ganze birationale Transformationen der Ebene*, J. Reine Angew. Math. 184, (1942), 161-174.

- [Ko] E.R. Kolchin, *Differential algebra and algebraic groups*, Pure and Applied Mathematics, Vol. 54, Academic Press New York-London, 1973.
- [KZ00] S. Kaliman and M. Zaidenberg, *Miyanishi's characterization of the affine 3-space does not hold in higher dimensions.*, Ann. Inst. Fourier, **50**(6), 1649–1669, 2000.
- [vdK] W. van der Kulk, *On polynomial rings in two variables*, Nieuw Arch. Wiskd., III. Ser. 1, 33–41 (1953).
- [ML1] L. Makar-Limanov, *On the hypersurface $x + x^2y + z^2 + t^3 = 0$ in $\mathbb{C}^4$ or a $\mathbb{C}^3$ -like threefold which is not $\mathbb{C}^3$* , Isr. J. Math. 96, Pt. B, 419–429 (1996).
- [ML2] L. Makar-Limanov, *Again $x + x^2y + z^2 + t^3 = 0$* , Affine algebraic geometry 177–182, Contemp. Math. **369**, Amer. Math. Soc., Providence, RI, 2005.
- [Sei] A. Seidenberg, *Derivations and integral closure*, Pacific J. Math. **16**, 1966, 167–173.
- [Sh] I. Shafarevich, *Basic algebraic geometry 1. Varieties in projective space*, second edition, Springer Verlag, Berlin, 1994.
- [SU] I.P. Shestakov and U.U. Umirbaev, *Poisson brackets and two-generated subalgebras of rings of polynomials*, J. Amer. Math. Soc. **17**(1), 2004, 181–196 (electronic).
- [SU2] I.P. Shestakov and U.U. Umirbaev, *The tame and the wild automorphisms of polynomial rings in three variables*, J. Amer. Math. Soc. **17**(1), 2004, 197–227 (electronic).
- [Vas] W.V. Vasconcelos, *Derivations of commutative noetherian rings*, Math. Z. **112**, 1969, 229–233.
- [V07] S. Vénéreau, *A parachute for the degree of a polynomial in algebraically independent ones*, arXiv:0704.1561, 2007.
- [Yu] J.T. Yu, *On relations between Jacobians and minimal polynomials*, Linear Algebra Appl. **221**, 1995, 19–29. MR **96c**:14014.